

Search for Fermion-Pair Decays $Q\bar{Q} \rightarrow (tW^\mp)(\bar{t}W^\pm)$ in Same-Charge Dilepton Events

T. Aaltonen,²⁶ J. Adelman,⁶⁵ B. Álvarez González^{v,13} S. Amerio^{dd,46} D. Amidei,³⁷ A. Anastassov,⁴¹ A. Annovi,²²
 J. Antos,¹⁷ G. Apollinari,²⁰ A. Apresyan,⁵¹ T. Arisawa,⁶¹ A. Artikov,¹⁸ J. Asaadi,⁵⁷ W. Ashmanskas,²⁰ A. Attal,⁴
 A. Aurisano,⁵⁷ F. Azfar,⁴⁵ W. Badgett,²⁰ A. Barbaro-Galtieri,³¹ V.E. Barnes,⁵¹ B.A. Barnett,²⁸ P. Barria^{ff,49}
 P. Bartos,¹⁷ G. Bauer,³⁵ P.-H. Beauchemin,³⁶ F. Bedeschi,⁴⁹ D. Beecher,³³ S. Behari,²⁸ G. Bellettini^{ee,49}
 J. Bellinger,⁶³ D. Benjamin,¹⁹ A. Beretvas,²⁰ D. Berge,¹⁶ A. Bhatti,⁵³ M. Binkley,²⁰ D. Bisello^{dd,46} I. Bizjak^{jj,33}
 R.E. Blair,² C. Blocker,⁷ B. Blumenfeld,²⁸ A. Bocci,¹⁹ A. Bodek,⁵² V. Boisvert,⁵² D. Bortoletto,⁵¹
 J. Boudreau,⁵⁰ A. Boveia,⁶⁶ B. Brau^{a,66} A. Bridgeman,²⁷ L. Brigliadori^{cc,6} C. Bromberg,³⁸ E. Brubaker,⁶⁵
 J. Budagov,¹⁸ H.S. Budd,⁵² S. Budd,²⁷ K. Burkett,²⁰ G. Busetto^{dd,46} P. Bussey,²⁴ A. Buzatu,³⁶ K. L. Byrum,²
 S. Cabrera^{x,19} C. Calancha,³⁴ S. Camarda,⁴ M. Campanelli,³³ M. Campbell,³⁷ F. Canelli^{14,20} A. Canepa,⁴⁸
 B. Carls,²⁷ D. Carlsmith,⁶³ R. Carosi,⁴⁹ S. Carrillo^{n,21} S. Carron,²⁰ B. Casal,¹³ M. Casarsa,²⁰ A. Castro^{cc,6}
 P. Catastini^{ff,49} D. Cauz,⁵⁸ V. Cavaliere^{ff,49} M. Cavalli-Sforza,⁴ A. Cerri,³¹ L. Cerrito^{q,33} S.H. Chang,³⁰
 Y.C. Chen,¹ M. Chertok,⁸ G. Chiarelli,⁴⁹ G. Chlachidze,²⁰ F. Chlebana,²⁰ K. Cho,³⁰ D. Chokheli,¹⁸ J.P. Chou,²⁵
 K. Chung^{o,20} W.H. Chung,⁶³ Y.S. Chung,⁵² T. Chwalek,²⁹ C.I. Ciobanu,⁴⁷ M.A. Ciocci^{ff,49} A. Clark,²³ D. Clark,⁷
 G. Compostella,⁴⁶ M.E. Convery,²⁰ J. Conway,⁸ M. Corbo,⁴⁷ M. Cordelli,²² C.A. Cox,⁸ D.J. Cox,⁸ F. Crescioli^{ee,49}
 C. Cuenca Almenar,⁶⁴ J. Cuevas^{v,13} R. Culbertson,²⁰ J.C. Cully,³⁷ D. Dagenhart,²⁰ M. Datta,²⁰ T. Davies,²⁴
 P. de Barbaro,⁵² S. De Cecco,⁵⁴ A. Deisher,³¹ G. De Lorenzo,⁴ M. Dell'Orso^{ee,49} C. Deluca,⁴ L. Demortier,⁵³
 J. Deng^{f,19} M. Deninno,⁶ M. d'Errico^{dd,46} A. Di Canto^{ee,49} G.P. di Giovanni,⁴⁷ B. Di Ruzza,⁴⁹ J.R. Dittmann,⁵
 M. D'Onofrio,⁴ S. Donati^{ee,49} P. Dong,²⁰ T. Dorigo,⁴⁶ S. Dube,⁵⁵ K. Ebina,⁶¹ A. Elagin,⁵⁷ R. Erbacher,⁸
 D. Errede,²⁷ S. Errede,²⁷ N. Ershaidat^{bb,47} R. Eusebi,⁵⁷ H.C. Fang,³¹ S. Farrington,⁴⁵ W.T. Fedorko,⁶⁵ R.G. Feild,⁶⁴
 M. Feindt,²⁹ J.P. Fernandez,³⁴ C. Ferrazza^{gg,49} R. Field,²¹ G. Flanagan^{s,51} R. Forrest,⁸ M.J. Frank,⁵ M. Franklin,²⁵
 J.C. Freeman,²⁰ I. Furic,²¹ M. Gallinaro,⁵³ J. Galyardt,¹⁴ F. Garbersen,⁶⁶ J.E. Garcia,²³ A.F. Garfinkel,⁵¹
 P. Garosi^{ff,49} H. Gerberich,²⁷ D. Gerdes,³⁷ A. Gessler,²⁹ S. Giagu^{hh,54} V. Giakoumopoulou,³ P. Giannetti,⁴⁹
 K. Gibson,⁵⁰ J.L. Gimmell,⁵² C.M. Ginsburg,²⁰ N. Giokaris,³ M. Giordani^{ii,58} P. Giromini,²² M. Giunta,⁴⁹
 G. Giurgiu,²⁸ V. Glagolev,¹⁸ D. Glenzinski,²⁰ M. Gold,⁴⁰ N. Goldschmidt,²¹ A. Golossanov,²⁰ G. Gomez,¹³
 G. Gomez-Ceballos,³⁵ M. Goncharov,³⁵ O. González,³⁴ I. Gorelov,⁴⁰ A.T. Goshaw,¹⁹ K. Goulianos,⁵³
 A. Gresele^{dd,46} S. Grinstein,⁴ C. Grosso-Pilcher,⁶⁵ R.C. Group,²⁰ U. Grundler,²⁷ J. Guimaraes da Costa,²⁵
 Z. Gunay-Unalan,³⁸ C. Haber,³¹ S.R. Hahn,²⁰ E. Halkiadakis,⁵⁵ B.-Y. Han,⁵² J.Y. Han,⁵² F. Happacher,²²
 K. Hara,⁵⁹ D. Hare,⁵⁵ M. Hare,⁶⁰ R.F. Harr,⁶² M. Hartz,⁵⁰ K. Hatakeyama,⁵ C. Hays,⁴⁵ M. Heck,²⁹ J. Heinrich,⁴⁸
 M. Herndon,⁶³ J. Heuser,²⁹ S. Hewamanage,⁵ M. Hickman,¹² D. Hidas,⁵⁵ C.S. Hill^{c,66} D. Hirschbuehl,²⁹
 A. Hocker,²⁰ S. Hou,¹ M. Houlden,³² S.-C. Hsu,³¹ R.E. Hughes,⁴² M. Hurwitz,⁶⁵ U. Husemann,⁶⁴ M. Hussein,³⁸
 J. Huston,³⁸ J. Incandela,⁶⁶ G. Introzzi,⁴⁹ M. Iori^{hh,54} A. Ivanov^{p,8} E. James,²⁰ D. Jang,¹⁴ B. Jayatilaka,¹⁹
 E.J. Jeon,³⁰ M.K. Jha,⁶ S. Jindariani,²⁰ W. Johnson,⁸ M. Jones,⁵¹ K.K. Joo,³⁰ S.Y. Jun,¹⁴ J.E. Jung,³⁰
 T.R. Junk,²⁰ T. Kamon,⁵⁷ D. Kar,²¹ P.E. Karchin,⁶² Y. Kato^{m,44} R. Kephart,²⁰ W. Ketchum,⁶⁵ J. Keung,⁴⁸
 V. Khotilovich,⁵⁷ B. Kilminster,²⁰ D.H. Kim,³⁰ H.S. Kim,³⁰ H.W. Kim,³⁰ J.E. Kim,³⁰ M.J. Kim,²² S.B. Kim,³⁰
 S.H. Kim,⁵⁹ Y.K. Kim,⁶⁵ N. Kimura,⁶¹ L. Kirsch,⁷ S. Klimenko,²¹ K. Kondo,⁶¹ D.J. Kong,³⁰ J. Konigsberg,²¹
 A. Korytov,²¹ A.V. Kotwal,¹⁹ M. Kreps,²⁹ J. Kroll,⁴⁸ D. Krop,⁶⁵ N. Krumnack,⁵ M. Kruse,¹⁹ V. Krutelyov,⁶⁶
 T. Kuhr,²⁹ N.P. Kulkarni,⁶² M. Kurata,⁵⁹ S. Kwang,⁶⁵ A.T. Laasanen,⁵¹ S. Lami,⁴⁹ S. Lammel,²⁰ M. Lancaster,³³
 R.L. Lander,⁸ K. Lannon^{u,42} A. Lath,⁵⁵ G. Latino^{ff,49} I. Lazzizzera^{dd,46} T. LeCompte,² E. Lee,⁵⁷ H.S. Lee,⁶⁵
 J.S. Lee,³⁰ S.W. Lee^{w,57} S. Leone,⁴⁹ J.D. Lewis,²⁰ C.-J. Lin,³¹ J. Linacre,⁴⁵ M. Lindgren,²⁰ E. Lipeles,⁴⁸ A. Lister,²³
 D.O. Litvintsev,²⁰ C. Liu,⁵⁰ T. Liu,²⁰ N.S. Lockyer,⁴⁸ A. Loginov,⁶⁴ L. Lovas,¹⁷ D. Lucchesi^{dd,46} J. Lueck,²⁹
 P. Lujan,³¹ P. Lukens,²⁰ G. Lungu,⁵³ J. Lys,³¹ R. Lysak,¹⁷ D. MacQueen,³⁶ R. Madrak,²⁰ K. Maeshima,²⁰
 K. Makhoul,³⁵ P. Maksimovic,²⁸ S. Malde,⁴⁵ S. Malik,³³ G. Manca^{e,32} A. Manousakis-Katsikakis,³ F. Margaroli,⁵¹
 C. Marino,²⁹ C.P. Marino,²⁷ A. Martin,⁶⁴ V. Martin^{k,24} M. Martínez,⁴ R. Martínez-Ballarín,³⁴ P. Mastrandrea,⁵⁴
 M. Mathis,²⁸ M.E. Mattson,⁶² P. Mazzanti,⁶ K.S. McFarland,⁵² P. McIntyre,⁵⁷ R. McNulty^{j,32} A. Mehta,³²
 P. Mehtala,²⁶ A. Menzione,⁴⁹ C. Mesropian,⁵³ T. Miao,²⁰ D. Mietlicki,³⁷ N. Miladinovic,⁷ R. Miller,³⁸ C. Mills,²⁵
 M. Milnik,²⁹ A. Mitra,¹ G. Mitselmakher,²¹ H. Miyake,⁵⁹ S. Moed,²⁵ N. Moggi,⁶ M.N. Mondragon^{n,20} C.S. Moon,³⁰
 R. Moore,²⁰ M.J. Morello,⁴⁹ J. Morlock,²⁹ P. Movilla Fernandez,²⁰ J. Mülmenstädt,³¹ A. Mukherjee,²⁰
 Th. Muller,²⁹ P. Murat,²⁰ M. Mussini^{cc,6} J. Nachtman^{o,20} Y. Nagai,⁵⁹ J. Naganoma,⁵⁹ K. Nakamura,⁵⁹
 I. Nakano,⁴³ A. Napier,⁶⁰ J. Nett,⁶³ C. Neu^{z,48} M.S. Neubauer,²⁷ S. Neubauer,²⁹ J. Nielsen^{g,31} L. Nodulman,²

M. Norman,¹⁰ O. Norriella,²⁷ E. Nurse,³³ L. Oakes,⁴⁵ S.H. Oh,¹⁹ Y.D. Oh,³⁰ I. Oksuzian,²¹ T. Okusawa,⁴⁴
 R. Orava,²⁶ K. Osterberg,²⁶ S. Pagan Griso^{dd},⁴⁶ C. Pagliarone,⁵⁸ E. Palencia,²⁰ V. Papadimitriou,²⁰
 A. Papaikonomou,²⁹ A.A. Paramanov,² B. Parks,⁴² S. Pashapour,³⁶ J. Patrick,²⁰ G. Paulettaⁱⁱ,⁵⁸ M. Paulini,¹⁴
 C. Paus,³⁵ T. Peiffer,²⁹ D.E. Pellett,⁸ A. Penzo,⁵⁸ T.J. Phillips,¹⁹ G. Piacentino,⁴⁹ E. Pianori,⁴⁸ L. Pinera,²¹
 K. Pitts,²⁷ C. Plager,⁹ L. Pondrom,⁶³ K. Potamianos,⁵¹ O. Poukhov*,¹⁸ F. Prokoshin^y,¹⁸ A. Pronko,²⁰
 F. Ptohosⁱ,²⁰ E. Pueschel,¹⁴ G. Punzi^{ee},⁴⁹ J. Pursley,⁶³ J. Rademacker^c,⁴⁵ A. Rahaman,⁵⁰ V. Ramakrishnan,⁶³
 N. Ranjan,⁵¹ I. Redondo,³⁴ P. Renton,⁴⁵ M. Renz,²⁹ M. Rescigno,⁵⁴ S. Richter,²⁹ F. Rimondi^{cc},⁶ L. Ristori,⁴⁹
 A. Robson,²⁴ T. Rodrigo,¹³ T. Rodriguez,⁴⁸ E. Rogers,²⁷ S. Rolli,⁶⁰ R. Roser,²⁰ M. Rossi,⁵⁸ R. Rossin,⁶⁶
 P. Roy,³⁶ A. Ruiz,¹³ J. Russ,¹⁴ V. Rusu,²⁰ B. Rutherford,²⁰ H. Saarikko,²⁶ A. Safonov,⁵⁷ W.K. Sakumoto,⁵²
 L. Santiⁱⁱ,⁵⁸ L. Sartori,⁴⁹ K. Sato,⁵⁹ A. Savoy-Navarro,⁴⁷ P. Schlabach,²⁰ A. Schmidt,²⁹ E.E. Schmidt,²⁰
 M.A. Schmidt,⁶⁵ M.P. Schmidt*,⁶⁴ M. Schmitt,⁴¹ T. Schwarz,⁸ L. Scodellaro,¹³ A. Scribano^{ff},⁴⁹ F. Scuri,⁴⁹
 A. Sedov,⁵¹ S. Seidel,⁴⁰ Y. Seiya,⁴⁴ A. Semenov,¹⁸ L. Sexton-Kennedy,²⁰ F. Sforza^{ee},⁴⁹ A. Sfyrla,²⁷ S.Z. Shalhout,⁶²
 T. Shears,³² P.F. Shepard,⁵⁰ M. Shimojima^t,⁵⁹ S. Shiraishi,⁶⁵ M. Shochet,⁶⁵ Y. Shon,⁶³ I. Shreyber,³⁹
 A. Simonenko,¹⁸ P. Sinervo,³⁶ A. Sisakyan,¹⁸ A.J. Slaughter,²⁰ J. Slaunwhite,⁴² K. Sliwa,⁶⁰ J.R. Smith,⁸
 F.D. Snider,²⁰ R. Snihur,³⁶ A. Soha,²⁰ S. Somalwar,⁵⁵ V. Sorin,⁴ P. Squillacioti^{ff},⁴⁹ M. Stanitzki,⁶⁴ R. St. Denis,²⁴
 B. Stelzer,³⁶ O. Stelzer-Chilton,³⁶ D. Stentz,⁴¹ J. Strologas,⁴⁰ G.L. Strycker,³⁷ J.S. Suh,³⁰ A. Sukhanov,²¹
 I. Suslov,¹⁸ A. Taffard^f,²⁷ R. Takashima,⁴³ Y. Takeuchi,⁵⁹ R. Tanaka,⁴³ J. Tang,⁶⁵ M. Tecchio,³⁷ P.K. Teng,¹
 J. Thom^h,²⁰ J. Thome,¹⁴ G.A. Thompson,²⁷ E. Thomson,⁴⁸ P. Tipton,⁶⁴ P. Ttito-Guzmán,³⁴ S. Tkaczyk,²⁰
 D. Toback,⁵⁷ S. Tokar,¹⁷ K. Tollefson,³⁸ T. Tomura,⁵⁹ D. Tonelli,²⁰ S. Torre,²² D. Torretta,²⁰ P. Totaroⁱⁱ,⁵⁸
 S. Tourneur,⁴⁷ M. Trovato^{gg},⁴⁹ S.-Y. Tsai,¹ Y. Tu,⁴⁸ N. Turini^{ff},⁴⁹ F. Ukegawa,⁵⁹ S. Uozumi,³⁰ N. van Remortel^b,²⁶
 A. Varganov,³⁷ E. Vataga^{gg},⁴⁹ F. Vázquezⁿ,²¹ G. Velev,²⁰ C. Vellidis,³ M. Vidal,³⁴ I. Vila,¹³ R. Vilar,¹³
 M. Vogel,⁴⁰ I. Volobouev^w,³¹ G. Volpi^{ee},⁴⁹ P. Wagner,⁴⁸ R.G. Wagner,² R.L. Wagner,²⁰ W. Wagner^{aa},²⁹
 J. Wagner-Kuhr,²⁹ T. Wakisaka,⁴⁴ R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³⁶ D. Waters,³³ M. Weinberger,⁵⁷
 J. Weinel,²⁹ W.C. Wester III,²⁰ B. Whitehouse,⁶⁰ D. Whiteson^f,⁴⁸ A.B. Wicklund,² E. Wicklund,²⁰ S. Wilbur,⁶⁵
 G. Williams,³⁶ H.H. Williams,⁶⁷ M.G. Wilson,⁵⁶ P. Wilson,²⁰ B.L. Winer,⁴² P. Wittich^h,²⁰ S. Wolbers,²⁰
 C. Wolfe,⁶⁵ H. Wolfe,⁴² T. Wright,³⁷ X. Wu,²³ F. Würthwein,¹⁰ A. Yagil,¹⁰ K. Yamamoto,⁴⁴ J. Yamaoka,¹⁹
 U.K. Yang^r,⁶⁵ Y.C. Yang,³⁰ W.M. Yao,³¹ G.P. Yeh,²⁰ K. Yi^o,²⁰ J. Yoh,²⁰ K. Yorita,⁶¹ T. Yoshida^l,⁴⁴ G.B. Yu,¹⁹
 I. Yu,³⁰ S.S. Yu,²⁰ J.C. Yun,²⁰ A. Zanetti,⁵⁸ Y. Zeng,¹⁹ X. Zhang,²⁷ Y. Zheng^d,⁹ and S. Zucchelli^{cc6}

(CDF Collaboration[†])

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^{cc}University of Bologna, I-40127 Bologna, Italy*

⁷*Brandeis University, Waltham, Massachusetts 02254*

⁸*University of California, Davis, Davis, California 95616*

⁹*University of California, Los Angeles, Los Angeles, California 90024*

¹⁰*University of California, San Diego, La Jolla, California 92093*

¹¹*University of California, Santa Barbara, Santa Barbara, California 93106*

¹²*University of California, Irvine, Irvine, California 92697*

¹³*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹⁴*Carnegie Mellon University, Pittsburgh, PA 15213*

¹⁵*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

¹⁶*European Organization for Nuclear Research, Geneva 23, Switzerland*

¹⁷*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹⁸*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁹*Duke University, Durham, North Carolina 27708*

²⁰*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

²¹*University of Florida, Gainesville, Florida 32611*

²²*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

²³*University of Geneva, CH-1211 Geneva 4, Switzerland*

²⁴*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²⁵*Harvard University, Cambridge, Massachusetts 02138*

²⁶*Division of High Energy Physics, Department of Physics,*

University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

- ²⁷University of Illinois, Urbana, Illinois 61801
- ²⁸The Johns Hopkins University, Baltimore, Maryland 21218
- ²⁹Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany
- ³⁰Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea
- ³¹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720
- ³²University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³³University College London, London WC1E 6BT, United Kingdom
- ³⁴Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
- ³⁵Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
- ³⁶Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³⁷University of Michigan, Ann Arbor, Michigan 48109
- ³⁸Michigan State University, East Lansing, Michigan 48824
- ³⁹Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ⁴⁰University of New Mexico, Albuquerque, New Mexico 87131
- ⁴¹Northwestern University, Evanston, Illinois 60208
- ⁴²The Ohio State University, Columbus, Ohio 43210
- ⁴³Okayama University, Okayama 700-8530, Japan
- ⁴⁴Osaka City University, Osaka 588, Japan
- ⁴⁵University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴⁶Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^{4d}University of Padova, I-35131 Padova, Italy
- ⁴⁷LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104
- ⁴⁹Istituto Nazionale di Fisica Nucleare Pisa, ^{4e}University of Pisa, ^{4f}University of Siena and ^{4g}Scuola Normale Superiore, I-56127 Pisa, Italy
- ⁵⁰University of Pittsburgh, Pittsburgh, Pennsylvania 15260
- ⁵¹Purdue University, West Lafayette, Indiana 47907
- ⁵²University of Rochester, Rochester, New York 14627
- ⁵³The Rockefeller University, New York, New York 10021
- ⁵⁴Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, ^{4h}Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵⁵Rutgers University, Piscataway, New Jersey 08855
- ⁵⁶SLAC National Accelerator Laboratory, Menlo Park, California 94025
- ⁵⁷Texas A&M University, College Station, Texas 77843
- ⁵⁸Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, ⁴ⁱUniversity of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁹University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁶⁰Tufts University, Medford, Massachusetts 02155
- ⁶¹Waseda University, Tokyo 169, Japan
- ⁶²Wayne State University, Detroit, Michigan 48201
- ⁶³University of Wisconsin, Madison, Wisconsin 53706
- ⁶⁴Yale University, New Haven, Connecticut 06520
- ⁶⁵Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637
- ⁶⁶University of California, Santa Barbara, Santa Barbara, California 93106
- ⁶⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104

We report the most restrictive direct limits on masses of fourth-generation down-type quarks b' , and quark-like composite fermions (B or $T_{5/3}$), decaying promptly to tW^\mp . We search for a significant excess of events with two same-charge leptons (e, μ), several hadronic jets, and missing transverse energy. An analysis of data from $p\bar{p}$ collisions with an integrated luminosity of 2.7 fb^{-1} collected with the CDF II detector at Fermilab yields no evidence for such a signal, setting mass limits $m_{b'}, m_B > 338 \text{ GeV}/c^2$ and $m_{T_{5/3}} > 365 \text{ GeV}/c^2$ at 95% confidence level.

PACS numbers: 12.60.-i, 13.85.Rm, 14.65.-q, 14.80.-j

*Deceased

†With visitors from ^aUniversity of Massachusetts Amherst,

The standard model (SM) of particle physics accommodates three generations of fundamental quarks and leptons, but does not prohibit a fourth. Recent measurements of charge-parity (CP) nonconservation in B -meson decays [1] have two features in common: they are more than two standard deviations from SM expectations, and they are sensitive to contributions [2] from a fourth-generation up-type quark, t' . This pattern of measurements [3–6], if genuine, warrants a search for another generation of quarks or a multiplet of quark-like objects. Another quark generation (Cf. [7]) could provide sources of particle-antiparticle asymmetries large enough to account for the baryon asymmetry of the universe [8], and accommodate a heavier Higgs boson (the source of electroweak symmetry breaking and mass generation) than a three-generation model [9].

This Letter reports a search for heavy particles Q decaying to a top quark and a W boson at a mass scale relevant to both the B -meson anomalies and the Higgs mechanism. We search for pair-production of $Q\bar{Q}$ via strong interactions, where Q is either a fourth-generation down-type quark b' or a quark-like (non-hadronic) composite fermion B or $T_{5/3}$ [10]. The B and $T_{5/3}$ (with $5/3$ electron charge) that we consider might arise from symmetries, consistent with precise electroweak measurements [11, 12]. If $T_{5/3}$ exists, the existence of B is implied, doubling the expected event rate. Many additional models of new phenomena that provide a Higgs mechanism also predict particles with large couplings to the third-generation t quark and similar decay modes. For instance, models of warped extra dimensions, equivalent to

models of strongly interacting composite particles, predict fermion excitations with the quantum numbers of quarks. A summary is given in [13].

In each case, $Q \rightarrow tW^\mp, t \rightarrow bW^+$ [14]. We investigate the case in which two same-charge W bosons decay leptonically (including τ decays to e or μ). This is the first search for quark-like particles in this mode [15], achieving the most sensitive direct limits on short-lived fourth-generation particles. (We do not consider long-lived particles, since displaced vertices require different analysis methods.)

We assume that Q decays exclusively to tW^\mp . This is expected for B and $T_{5/3}$, and for b' it is expected [16] under the assumptions that (a) coupling to light quarks is insignificant, (b) $m_{b'} > m_{top} + m_W = 255 \text{ GeV}/c^2$, and (c) $|m_{t'} - m_{b'}| < m_W$. These assumptions are justified by experimental constraints. A search for $Q \rightarrow W + \text{jet}$ [17] found $m_{t'} > 311 \text{ GeV}/c^2$, implying a similar limit on $m_{b'}$ if the b' decay to this channel is significant. Combining this limit with results of a search for $Q \rightarrow Z + \text{jets}$ [18] and an analysis of branching fractions for b' [16], we infer $m_{b'} > 255 \text{ GeV}/c^2$. A fourth generation is most consistent with precise electroweak measurements when the mass splitting Δm between b' and t' is less than the W -boson mass but non-zero; ref. [9] gives $\Delta m \approx 50 \text{ GeV}/c^2$, based on $|m_{t'} - m_{b'}| < m_W$.

We use a data sample corresponding to an integrated luminosity of 2.7 fb^{-1} collected with the CDF II detector [19] at the Tevatron $p\bar{p}$ collider at Fermilab. The data acquisition system is triggered by e or μ candidates with $p_T > 18 \text{ GeV}/c$ [20]. We require the $\ell^\pm \ell^\pm b j \cancel{E}_T$ signature, following [21]: two same-charge reconstructed leptons (e or μ) with pseudorapidity magnitude $|\eta| < 1.1$ and $p_T > 20 \text{ GeV}/c$, where at least one lepton is isolated [22]; at least two jets with $E_T > 15 \text{ GeV}$ and $|\eta| < 2.4$; at least one of the jets with evidence of a long-lived particle (b -tag) using the tight SECVTX algorithm [23]; and missing transverse energy $\cancel{E}_T > 20 \text{ GeV}$ [24].

The dominant background comes from events in which one of the leptons is a misidentified light-flavor jet or a lepton from the decay of a bottom or charmed hadron in a heavy flavor jet, largely from W production in association with light or heavy flavor jets or from $t\bar{t}$ production with semi-leptonic decays. This background is described using a lepton misidentification model from inclusive jet data [25] applied to $W + \text{jet}$ events. In same-charge dilepton control regions without a b -tag requirement, this model describes well the kinematics of observed events with large missing transverse energy. Nevertheless, the requirement of a b -tag in the final selection introduces uncertainty regarding the misidentification model, leading to a final 100% systematic uncertainty, as described in [21].

Other backgrounds include processes that produce electron-positron pairs. These may be reconstructed with the same charge due to asymmetric γ conversions in the

Amherst, Massachusetts 01003, ^bUniversiteit Antwerpen, B-2610 Antwerp, Belgium, ^cUniversity of Bristol, Bristol BS8 1TL, United Kingdom, ^dChinese Academy of Sciences, Beijing 100864, China, ^eIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^fUniversity of California Irvine, Irvine, CA 92697, ^gUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^hCornell University, Ithaca, NY 14853, ⁱUniversity of Cyprus, Nicosia CY-1678, Cyprus, ^jUniversity College Dublin, Dublin 4, Ireland, ^kUniversity of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, ^lUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017 ^mKinki University, Higashi-Osaka City, Japan 577-8502 ⁿUniversidad Iberoamericana, Mexico D.F., Mexico, ^oUniversity of Iowa, Iowa City, IA 52242, ^pKansas State University, Manhattan, KS 66506 ^qQueen Mary, University of London, London, E1 4NS, England, ^rUniversity of Manchester, Manchester M13 9PL, England, ^sMuons, Inc., Batavia, IL 60510, ^tNagasaki Institute of Applied Science, Nagasaki, Japan, ^uUniversity of Notre Dame, Notre Dame, IN 46556, ^vUniversity of Oviedo, E-33007 Oviedo, Spain, ^wTexas Tech University, Lubbock, TX 79609, ^xIFIC(CSIC)Universitat de Valencia), 56071 Valencia, Spain, ^yUniversidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, ^zUniversity of Virginia, Charlottesville, VA 22906 ^{aa}Bergische Universität Wuppertal, 42097 Wuppertal, Germany, ^{bb}Yarmouk University, Irbid 211-63, Jordan ^{jj}On leave from J. Stefan Institute, Ljubljana, Slovenia,




FIG. 1: Missing transverse energy in events with same-charge leptons in 2.7 fb^{-1} . The right outer-most bin includes overflow events with $E_T > 160 \text{ GeV}$.

process $e_{\text{hard}}^- \rightarrow e_{\text{soft}}^- \gamma \rightarrow e_{\text{hard}}^+ e_{\text{soft}}^- e_{\text{soft}}^-$, where hard and soft refer to large and small transverse momentum, respectively. The major contributions from this mechanism are from events with a Z or virtual γ in association with jets ($Z/\gamma^* + \text{jets}$) and $t\bar{t}$ production with fully leptonic decays.

Estimates of the backgrounds from $Z/\gamma^* + \text{jets}$ processes are made with the ALPGEN [26] v2.10 simulation code interfaced with PYTHIA 6.325 [27] in the MLM scheme [26] for the hadronization and fragmentation and normalized to data in opposite-charge events in the Z mass region. The detector response for both $Z + \text{jets}$ and $t\bar{t}$ processes is evaluated using the CDF simulation program CDFSIM [28], where, to avoid double-counting, the same-charge leptons are required to originate from the W or Z decays rather than from misidentified jets.

To validate the modeling of the rate of hard bremsstrahlung from electrons, we compare our prediction for the contribution of $Z \rightarrow e^+e^-$ to the observed sample of same-charge electrons or positrons without a b -tag or missing transverse energy requirement. The shape of the dilepton invariant mass spectrum and yield in the Z mass region ($M_{ll} = [M_Z - 20, M_Z + 20]$) agrees well with the prediction. In addition, $\mu\mu$ and $e\mu$ events have negligible contributions from hard bremsstrahlung, as predicted. Figure 1 shows that the missing transverse

TABLE I: Expected background contributions to the ee , $e\mu$, and $\mu\mu$ channels in 2.7 fb^{-1} from (a) Z and diboson, (b) $t\bar{t} \rightarrow \ell^+ \nu b \ell^- \nu \bar{b}$, and (c) misidentified lepton.

Source	ee	$\mu\mu$	$e\mu$	Total $\ell\ell$
(a)	0.01 ± 0.01	0	0.02 ± 0.02	0.03 ± 0.03
(b)	0.06 ± 0.04	0	0.09 ± 0.03	0.15 ± 0.05
(c)	0.6 ± 0.6	0.3 ± 0.3	0.5 ± 0.5	1.4 ± 1.4
Total	0.7 ± 0.6	0.3 ± 0.3	0.6 ± 0.5	1.6 ± 1.4
Data	0	1	1	2




FIG. 2: Number of reconstructed jets for the expected backgrounds. The observed data and the b' (or B) signal are shown at the best-fit rate for $m_Q = 330 \text{ GeV}/c^2$. The fitted size and shape for the $T_{5/3} + B$ signal is nearly identical.

energy in inclusive same-charge dilepton events is well described.

The $t\bar{t} \rightarrow \ell^+ \nu b \ell^- \nu \bar{b}$ backgrounds are estimated using events generated in PYTHIA 6.216 at $m_t = 172.5 \text{ GeV}/c^2$, assuming a $t\bar{t}$ production cross-section of 7.2 pb . Modeling of the $t\bar{t}$ contribution is validated by comparing predicted and observed rates of events with opposite-charge leptons, large E_T , and at least one b -tagged jet, where $t\bar{t}$ is expected to dominate.

Backgrounds to the $\ell^\pm \ell^\pm b j E_T$ signature with real same-charge leptons are rare in the SM; they are largely from WZ and ZZ production and are highly suppressed by the requirement of a b -tag. Backgrounds from diboson production WW , WZ , ZZ , $W\gamma$, and $Z\gamma$ in association with b jets are modeled with PYTHIA 6.216 and BAUR [29] generators.

Backgrounds from charge mismeasurement are insignificant, as the charge of a particle with $p_T \approx 100 \text{ GeV}/c$ is typically determined with more than 5σ significance [30]. Charge mismeasurement is very rare in this range, confirmed by the absence of any strong fea-

TABLE II: Theoretical cross-sections (σ_{NLO} in fb [36, 37]), expected yield (N), median expected 95% C.L limit ($\sigma_{\text{exp'd}}$ in fb), and observed 95% C.L limit (σ_{obs} in fb) for b' (or B) and ($T_{5/3} + B$) signals at varying masses.

Mass [GeV/c^2]	300	310	320	330	340	350	375	400
$b' \text{ or } B$	σ_{NLO}	227	176	137	106	83	64	34
	N	13.4	9.6	7.5	5.9	4.6	3.5	1.9
	$\sigma_{\text{exp'd}}$	67	63	63	62	63	63	63
	σ_{obs}	67	96	83	94	85	83	78
$T_{5/3} + B$	σ_{NLO}	454	352	274	212	166	128	68
	N	27.0	19.5	15.3	11.9	9.4	7.1	3.6
	$\sigma_{\text{exp'd}}$	86	89	69	62	59	65	66
	σ_{obs}	86	89	69	98	91	83	79




FIG. 3: Event displays for the observed three-jet, $\mu\mu$ event (a,b) and the five-jet, $e\mu$ event (c,d). Shown in (a) and (c) are views of the events along the beam axis; jets shown as cones, electrons as solid lines, muons as dotted lines and missing transverse energy as an arrow; lengths are proportional to p_T (see Table III). Shown in (b) and (d) are views of the events in $\eta - \phi$; jets shown as open circles, electrons as filled circles and muons as dashed circles; radii are proportional to p_T .

tures in dilepton invariant mass in the Z mass region in same-charge muon events. The largest potential source comes from $t\bar{t}$ events, in which the lepton momenta are typically smaller than $100 \text{ GeV}/c$. The final background estimates are given in Table I.

The b' and $T_{5/3} + B$ signals are modeled with the MADGRAPH simulation program following the minimal composite Higgs model described in [10] and paired with PYTHIA for hadronization and fragmentation. The acceptance is approximately 2.2%, nearly independent of heavy quark masses in the range $300\text{--}400 \text{ GeV}/c^2$. The expected numbers of events for b' (or B), and $T_{5/3} + B$ are given in Table II.

We observe two events in the signal region, in agreement with the expected backgrounds (see Table I). To calculate the most likely signal cross section, we perform a binned maximum-likelihood fit to the number of reconstructed jets. Figure 2 shows the number of recon-

TABLE III: Transverse momentum (in GeV/c) of leptons and transverse energy (in GeV) of jets in the two events with the $\ell^\pm \ell^\pm b j \cancel{E}_T$ signature.

Event	ℓ_1	ℓ_2	jet ₁	b-jet	\cancel{E}_T	other jets
$\mu^+ \mu^+$	80	31	78	25	87	40
$e^+ \mu^+$	73	21	60	42	27	39, 33, 24




FIG. 4: Theoretical cross sections for b' (or B) and $T_{5/3} + B$ with expected and observed 95% C.L. limits overlaid.

structed jets in the observed events, as well as the signal distribution with the best-fit value of the signal cross section. Kinematics of the two signal events is shown in Fig. 3 and the p_T values are given in Table III.

We construct confidence intervals [31] in the theoretical cross section by generating ensembles of simulated experiments that describe expected fluctuations of statistical and systematic uncertainties, including uncertainties in the jet-energy scale [32], gluon radiation [33], signal and background normalization, and parton distribution functions [34, 35]. The median expected and observed limits along with the theoretical next-to-leading-order (NLO) cross section [36, 37] are given in Table II and shown in Fig. 4.

We convert limits on the pair-production cross sections to limits on the fermion masses and obtain $m_{b'}, m_B > 338 \text{ GeV}/c^2$ and $m_{T_{5/3}} > 365 \text{ GeV}/c^2$ at 95% confidence level. The two events observed are consistent with the predicted number of background events, although we note that the $e\mu$ event has a number of jets characteristic of the signal, reducing the observed lower limits from what is expected. This is the most restrictive direct lower limit on the mass of a down-type fourth-generation quark, significantly reducing the allowed SM mass range, and the first lower limits on the masses of an exotic doublet $T_{5/3} + B$, which may figure prominently in future searches.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the

Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

-
- [1] Specifically, we refer to the mixing-induced CP asymmetry in the decays $B_s \rightarrow J/\psi \phi$ [3], the difference between direct CP asymmetries in the decays $B^0 \rightarrow K^+ \pi^-$ and $B^+ \rightarrow K^+ \pi^0$ [4, 6], and the values of mixing-induced CP asymmetry obtained from $B^0 \rightarrow J/\psi K_S^0$ or $B^0 \rightarrow (\phi, \eta', K_S^0 K_S^0) K_S^0$ [5, 6].
- [2] W.-S. Hou, M. Nagashima, and A. Soddu, Phys. Rev. Lett. **95**, 141601 (2005); W.-S. Hou, M. Nagashima, G. Raz, and A. Soddu, J. High Energy Phys. 09 (2006) 012; W.-S. Hou, H. n. Li, S. Mishima, and M. Nagashima, Phys. Rev. Lett. **98**, 131801 (2007); W.-S. Hou, M. Nagashima, and A. Soddu, Phys. Rev. D **76**, 016004 (2007).
- [3] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **100**, 161802 (2008); V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **101**, 241801 (2008); T. Aaltonen *et al.* (CDF Collaboration), CDF Public Note CDF9458, Aug. 7, 2008.
- [4] S.-W. Lin, Y. Unno, W.-S. Hou, and P. Chang *et al.* (Belle Collaboration), Nature **452**, 332 (2008); B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **76**, 091102 (2007); B. Aubert *et al.* (BABAR Collaboration), arXiv:hep-ex/0807.4226 [hep-ex].
- [5] E. Lunghi and A. Soni, Phys. Lett. B **666**, 162 (2008).
- [6] E. Barberio *et al.* (Heavy Flavor Averaging Group), arXiv:hep-ex/0808.1297v3 [hep-ex].
- [7] P. H. Frampton, P. Q. Hung, and M. Sher, Phys. Rept. **330**, 263 (2000).
- [8] W.-S. Hou, Chin. J. Phys. **47**, 134 (2009).
- [9] G. D. Kribs, T. Plehn, M. Spannowsky, and T. M. P. Tait, Phys. Rev. D **76**, 075016 (2007).
- [10] R. Contino and G. Servant, J. High Energy Phys. 06 (2008) 026.
- [11] P. Sikivie, L. Susskind, M. B. Voloshin, and V. I. Zakharov, Nucl. Phys. **B173**, 189 (1980).
- [12] K. Agashe, R. Contino, L. Da Rold, and A. Pomarol, Phys. Lett. B **641**, 62 (2006).
- [13] R. Contino, T. Kramer, M. Son, and R. Sundrum, J. High Energy Phys. 05 (2007) 074.
- [14] Unless otherwise indicated, particle types and decay processes imply also their charge conjugates.
- [15] Sensitivity studies have previously been shown by Ref [10] and the CMS Collaboration, CMS Physics Analysis Summary EXO-08-009, Aug. 29, 2008.
- [16] P. Q. Hung and M. Sher, Phys. Rev. D **77**, 037302 (2008).
- [17] A. Lister (for the CDF Collaboration), arXiv:hep-ex/0810.3349.
- [18] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **76**, 072006 (2007).
- [19] D. E. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [20] CDF uses a cylindrical coordinate system with the z axis along the proton beam axis. Pseudorapidity is $\eta \equiv -\ln(\tan(\theta/2))$, where θ is the polar angle relative to the proton beam direction, and ϕ is the azimuthal angle while $p_T = |p| \sin \theta$, $E_T = E \sin \theta$.
- [21] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **102**, 041801 (2009).
- [22] A lepton is isolated if the calorimeter energy in a cone $\Delta R < 0.4$ surrounding the lepton is less than 10% of the energy of the lepton.
- [23] A. Abulencia *et al.*, Phys. Rev. D **74**, 072006 (2006)
- [24] Missing transverse energy, \cancel{E}_T , is defined as the magnitude of the vector $-\sum_i E_T^i \vec{n}_i$ where E_T^i are the magnitudes of transverse energy contained in each calorimeter tower i , and \vec{n}_i is the unit vector from the interaction vertex to the tower in the transverse (x, y) plane. \cancel{E}_T is further corrected for the energy of identified muons.
- [25] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **93**, 142001 (2004).
- [26] M.L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. Polosa, J. High Energy Phys. 07 (2003) 001.
- [27] T. Sjostrand *et al.*, Comput. Phys. Commun. **238** 135 (2001).
- [28] T. Affolder *et al.* (CDF Collaboration), Nucl. Instrum. Methods **447**, 1 (2000).
- [29] U. Baur and E.L. Berger, Phys. Rev. D **41**, 1476 (1990).
- [30] A. Abulencia *et al.* (CDF Collaboration), J. Phys. G: Nucl. Part. Phys. **34** (2007) 2457.
- [31] G. J. Feldman and R. D. Cousins, Phys. Rev. D **57**, 3873 (1998).
- [32] A. Bhatti *et al.*, Nucl. Instrum. Methods **566**, 375 (2006).
- [33] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D. **73** 32003 (2006).
- [34] J. Pumplin *et al.* (CTEQ Collaboration), Journ. High. En. Phys. 07 (2002) 012.
- [35] A. D. Martin *et al.* (MRST Collaboration), Phys. Lett. B **356** 89 (1995).
- [36] R. Bonciani, S. Catani, M. L. Mangano, and P. Nason, Nucl. Phys. **B529**, 424 (1998).
- [37] M. Cacciari, S. Frixione, M. L. Mangano, P. Nason, and G. Ridolfi, J. High Energy Phys. 04 (2004) 068.