Properties of heavy quark hadrons at CDF

Juan Pablo Fernández Ramos

C.I.E.M.A.T.

29/04/2009

(for the CDF Collaboration)

Properties:

- Focus on masses , lifetimes (decays)

Why study B hadrons ?

- Heavy quark hadrons are the hydrogen atom of QCD => study of B hadron states = study of (non-perturbative) QCD
- Measurements of B hadron masses provide sensitive tests of potential models, HQET and all aspects of QCD including lattice gauge calculations

Why B Hadron Lifetimes ?

- Spectator model : b quark decays like free particle
=> all B hadron lifetimes are equal
- In reality QCD => lifetimes of B hadrons study the interplay between strong and weak interaction

Introduction to the Tevatron



- ppbar collisions at 1.96 TeV
- Excellent performance of Tevatron accelerator
- Keep breaking record of peak Initial Luminosity after 8 years of running
- CDF and D0 have already > **5 fb^{-1}** on tape
- Expect 6 fb^{-1} by end 2009 (8 after 2010 extension)
- The analyses presented in this talk span from 1 to 3.8 fb^{-1}

CDF detector





- Drift chamber (COT)
  ⇒ Good tracking resolution
  $\sigma(p_T)/p_T \sim 0.1\ \%\ \mathrm{GeV}^{-1}$
- Silicon vertex detector
  ⇒ Good vertex resolution
  ⇒ Important for triggering
  (using a displaced track trigger, SVT)
- TOF detector and dE/dx from COT
  ⇒ Good particle identification
- Muon System up to $|\eta| < 1.5$
  ⇒ Important for triggering

Basics of B Physics at the Tevatron









- High cross section σ ($\mathbf{p\bar{p} \rightarrow b\bar{b}}$) ~ 40 μb at $\sqrt{s}$ = 2 TeV
- Quarks fragment into hadrons: B_c^- ($b\bar{c}$), Λ_b(bdu), Σ_b^+ (buu), Σ_b^- (bdd) [Tevatron exclusive], B_s^0 ($b\bar{s}$), $\bar{B}_0$($b\bar{d}$), B^-($b\bar{u}$), also B*, B**, etc
- → Tevatron can be considered as a B factory

Online B selection process

- Huge background to the process $\sigma(\mathbf{p\bar{p}} \rightarrow \mathbf{b\bar{b}})$ in Tevatron: O(0.05 b)!
- B hadrons are filtered online using selective triggers based on clear signatures that overcome the QCD background :
  - events selected by a $J/\psi \rightarrow \mu\mu$ oriented dimuon trigger
  - events selected by an impact parameter based trigger (SVT)





Results :

- B_c mass, lifetime and cross section
- Λ_b lifetime in $\Lambda_b \rightarrow \Lambda_c \pi$
- B_s lifetime
- Σ_b Baryon

B_c lifetime



- B_c contains two heavy quarks: bottom and anti-charm
- Produced only at the Tevatron → unique testing ground for QCD
- Lifetime measurement in semileptonic decays B_c → J/Ψ+ l+X

$$\tau c\,(B_c) = 142.5\ ^{+15.8}_{-14.8}(\text{stat.}) \pm 5.5\ (\text{syst.})\ \mu\text{m}$$

$$\tau = 0.475\ ^{+0.053}_{-0.049}(\text{stat.}) \pm 0.018\ (\text{syst.})\ \text{ps}$$

Theoretical predictions:
0.47-0.59 ps



http://www-cdf.fnal.gov/physics/new/bottom/080327.blessed-BC_LT_SemiLeptonic/

B_c mass

- Best mass measurement from CDF in $B_c \rightarrow J/\Psi\pi$ decays (2.4 fb^{-1})

  $m(B_c) = 6275.6 \pm 2.9 \pm 2.5$ MeV/c (PRL 100:182002,2008)

good agreement with theory (6304 ± 12+18-0) Phys. Rev. Lett. 94, 172001 (2005)





Horizontal axis corresponds to different theory predictions

Experimental measurements with small uncertainties start to challenge theoretical models and lattice techniques

- Recent update (4.7 fb^{-1})
  - analysis framework is unchanged from PRL
  - uses improved track parameter uncertainties

B_c cross section

- Ratio of the production cross section times branching fraction of $B_c^+ \rightarrow J/\Psi\ \mu\ \nu$ relative to $B^+ \rightarrow J/\Psi K^+$ for two $p_T(B)$ cuts with 1 fb^{-1} .
- Select sample of 229(214) events with 3rd muon for $p_T(B_c)>4GeV/c$ (6)
- Estimate background contributions using data and PYTHIA samples
  - misidentified J/Ψ
  - misidentified muons
  - $b\bar{b}$ background
  - other b hadrons
- Total background : 111 ± 8 (107 ± 8) events.
- Select $B^+ \rightarrow J/\Psi K^+$
- Estimate the ratio



http://www-cdf.fnal.gov/physics/new/bottom/090305.blessed-bc-slXsec/

B_c cross section

$$\frac{\sigma(B_c^+)\cdot BR(B_c^+ \to J/\psi + \mu^+ + \nu)}{\sigma(B^+)\cdot BR(B^+ \to J/\psi + K^+)} = \frac{N_{B_c^+}}{N_{B^+}} \times \epsilon_{rel}$$

- ε_{rel} , relative efficiency :

  $\varepsilon_{rel} = \varepsilon_{B+} / \varepsilon_{Bc}$.

Efficiencies	Pt(B) > 4 GeV/c	Pt(B) > 6 GeV/c
$\varepsilon_{B+c}(\%)$	0.0551	0.1232
$\varepsilon_{B+}(\%)$	0.3231	0.6005
ε_{rel}	5.867 ± 0.068	4.873 ± 0.060

	Pt (B) > 4 GeV/c	Pt(B) > 6 GeV/c
$N(B_c^+)$	$117.6 \pm 17.2_{-6.4}^{+8.3}$	$107.1 \pm 16.7_{-6.1}^{+7.9}$
$N(B^+)$	2333 ± 55	2299 ± 53
ε_{rel}	5.867 ± 0.068	4.873 ± 0.060
	$-0.450^{+0.554} \pm 0.720$	$-0.278^{+0.420} \pm 0.298$
$\frac{N(B_c^+)}{N(B^+)} \times \varepsilon_{rel}$	0.295 ± 0.040 (stat)	0.227 ± 0.033 (stat)
	$-0.026^{+0.033}$(sys) ± 0.036(spectrum)	$-0.017^{+0.024}$(sys)± 0.014

Λ_b lifetime in $\Lambda_b \to \Lambda_c \pi$



- Important test of models that describe interaction between heavy and light quarks within bound states
- Precise theoretical predictions difficult due to QCD effects
- OPE/HQET predicts lifetime hierarchy of the b-hadrons
- CDF analysis with large sample of ~3000 signal events in 1.1 fb^{-1}
- Displaced track trigger requirements : 120 μm< IP < 1 mm
- Trigger bias corrected using simulation
- Sample composition obtained from mass distribution



Λ_b lifetime results

- PDF is convolution of : exponential , $c\tau$ resolution PDF, trigger eff.
- Most precise measurement: $\tau c(\Lambda_b) = 420.1 \pm 13.7(stat.) \pm 10.6\ (syst.)\ \mu m$
- $\tau(\Lambda_b)/\tau(B_0) = 0.916 \pm 0.038$
- Good agreement with theory (0.88 ± 0.05) and previous world average



arxiv:hep-ph:0310241



B_s lifetime in $B_s \rightarrow J/\Psi\phi$

- As part of angular analysis that together with three polarization amplitudes meaures $\Delta\Gamma$ and $\sin 2\beta_s$
- No bias from trigger
- Mean lifetime (assuming no CP violation, $\beta_s^{J/\Psi\phi} = 0$) :
  - $\tau c\ (Bs) = 1/\Gamma_s = 459 \pm 12$ (stat.) $\pm$ 3 (syst.) μm

    Results are mean of heavy and light states

    No flavor specific measurement



3200 candidates with no PID used in selection



http://www-cdf.fnal.gov/physics/new/bottom/080724.blessed-tagged_BsJPsiPhi_update_prelim/

B_s lifetime summary

- New measurement greater than earlier average
  - Strongly weighted by semileptonic channel from D0
- New combination will reduce discrepancy between data and expectation $\tau(B_s) \approx \tau(B^0)$



Σ_b , Σ_b^* Baryons

- Until 2006 Λ_b was only established B baryon

=> search for $\Sigma_b^+ = |buu\rangle$ $\Sigma_b^- = |bdd\rangle$

- $\Sigma_b^{*\pm}$ excited state of $\Sigma_b^{\pm}$ with s = 3/2
- Use two different triggers :
  - displaced track [PRL 99, 202001(2007);

    decay $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$ with $\Lambda_c^+ \rightarrow pK^-\pi^+$]
  - J/Ψ trigger [$\Lambda_b \rightarrow$ J/Ψ Λ]:

Reconstruct ~ 1600 Λ_b in J/Ψ Λ decay with 3.8 fb^{-1} .





Σ_b , Σ_b^* Baryons

State	Yield	Q or $\Delta_{\Sigma_b^*}$ (MeV/c^2)	Mass (MeV/c^2)
Σ_b^+	32^{+13+5}_{-12-3}	$Q_{\Sigma_b^+} = 48.5^{+2.0+0.2}_{-2.2-0.3}$	$5807.8^{+2.0}_{-2.2} \pm 1.7$
Σ_b^-	59^{+15+9}_{-14-4}	$Q_{\Sigma_b^-} = 55.9 \pm 1.0 \pm 0.2$	$5815.2 \pm 1.0 \pm 1.7$
Σ_b^{*+}	77^{+17+10}_{-16-6}	$\Delta_{\Sigma_b^*} = 21.2^{+2.0+0.4}_{-1.9-0.3}$	$5829.0^{+1.6+1.7}_{-1.8-1.8}$
Σ_b^{*-}	69^{+18+16}_{-17-5}		$5836.4 \pm 2.0^{+1.8}_{-1.7}$

- Q for $\Lambda_b \to \Lambda_c \pi$ decay mode



- Q for $\Lambda_b \to J/\Psi\ \Lambda$ decay mode



- Observe peaks with > 5σ w.r.t no signal
- Data suggest that in $\Lambda_b \to J/\Psi\ \Lambda$ decay mode there are consistent peaks

Conclusions

Very rich heavy flavour program at CDF.

Many results on properties of heavy B hadrons :

- B_c lifetime, mass and cross section
- Λ_b lifetime
- B_s lifetime
- heavy baryons Σ_b, Σ_b^* established

Great Tevatron performance.

More data to come.

Good time for flavor physics at the Tevatron right now!

Back up